

Checkpoint Task: Enthalpy changes

Learner Activity

Exothermic reactions

1. Write a definition of an exothermic reaction.

2. Draw an enthalpy profile diagram for an exothermic reaction.
Label the axes, ΔH and the activation energy.



3. Give an example of an exothermic reaction.

Endothermic reactions

4. Write a definition of an endothermic reaction.

5. Draw an enthalpy profile diagram for an endothermic reaction.

Label the axes, ΔH and the activation energy.



6. Give an example of an endothermic reaction.

Bond enthalpy

7. Write a definition of bond enthalpy. (You might know this term as 'bond energy'.)

8. In a chemical reaction, bonds in the reactants are broken, and new bonds are formed to make the products. Complete the following sentences.

Energy is to break bonds.

Energy is when bonds are formed.

The overall energy change of a reaction is the

Calculations


Remember:

enthalpy change = energy required to break bonds – energy released in making bonds

or

$$\Delta_r H = \Sigma(\text{bond enthalpies in reactants}) - \Sigma(\text{bond enthalpies in products})$$

9. Use bond enthalpies to calculate the enthalpy change for the following reaction.





Bond	H–H	Br–Br	H–Br
Bond enthalpy / kJ mol ⁻¹	438	193	366

Energy required to break bonds:

Energy released in forming new bonds:

Enthalpy change:

10. Use bond enthalpies to calculate the enthalpy change for the combustion of methane.



Bond	C-H	C-C	O-H	C=O	O=O
Bond enthalpy / kJ mol^{-1}	413	347	464	805	498

Energy required to break bonds:

Energy released in forming new bonds:


Enthalpy change:

11.


Bond	C-H	C-C	H-H	C=C
Bond enthalpy / kJ mol ⁻¹	413	347	436	612

Use the bond energies above to calculate

a) the enthalpy change for the hydrogenation of ethene



b) the enthalpy change for the cracking of decane



12. Explain in terms of bond breaking and bond formation why combustion reactions are exothermic but cracking reactions are endothermic.


Exothermic reactions

1. Write a definition of an exothermic reaction.

An exothermic reaction is one in which energy is given out to the surroundings.

2. Draw an enthalpy profile diagram for an exothermic reaction.

Label the axes, ΔH and the activation energy.



CHEMISTRY A AND CHEMISTRY B (SALTERS)**Learner Activity**

3. Give an example of an exothermic reaction.

E.g. combustion, lots of oxidation reactions (including rusting), neutralisation.


Endothermic reactions

4. Write a definition of an endothermic reaction.

An endothermic reaction is one in which energy is taken in from the surroundings.

5. Draw an enthalpy profile diagram for an endothermic reaction.

Label the axes, ΔH and the activation energy.



6. Give an example of an endothermic reaction.

E.g. decomposition of metal carbonates, electrolysis, reaction between ethanoic acid and sodium carbonate, cracking, photosynthesis.

Bond enthalpy

7. Write a definition of bond enthalpy. (You might know this term as 'bond energy'.)

The energy needed to break one mole of a given bond in a gaseous molecule. The units are kJ mol^{-1} (kilojoules per mole).

8. In a chemical reaction, bonds in the reactants are broken, and new bonds are formed to make the products. Complete the following sentences.


Energy is to break bonds.

Energy is when bonds are formed.

The overall energy change of a reaction is the

CHEMISTRY A AND CHEMISTRY B (SALTERS)**Learner Activity****Calculations**

9. Use bond enthalpies to calculate the enthalpy change for the following reaction.



Bond	H–H	Br–Br	H–Br
Bond enthalpy / kJ mol^{-1}	438	193	366

Energy required to break bonds:

$$(\text{H–H}) + (\text{Br–Br}) = 631 \text{ kJ mol}^{-1}$$



Energy released in forming new bonds:

$$2 \times (\text{H–Br}) = 732 \text{ kJ mol}^{-1}$$

Enthalpy change:

$$631 - 732 = -101 \text{ kJ mol}^{-1}$$

10. Use bond enthalpies to calculate the enthalpy change for the combustion of methane.



Bond	C–H	C–C	O–H	C=O	O=O
Bond enthalpy / kJ mol^{-1}	413	347	464	805	498

Energy required to break bonds:

$$4 \times (\text{C–H}) + 2 \times (\text{O=O}) = 2648 \text{ kJ mol}^{-1}$$

Energy released in forming new bonds:

$$2 \times (\text{C=O}) + 4 \times (\text{O–H}) = 3466 \text{ kJ mol}^{-1}$$

Enthalpy change:

$$2648 - 3466 = -818 \text{ kJ mol}^{-1}$$


CHEMISTRY A AND CHEMISTRY B (SALTERS)**Learner Activity**

11.

Bond	C–H	C–C	H–H	C=C
Bond enthalpy / kJ mol ⁻¹	413	347	436	612


Use the bond energies above to calculate

a) the enthalpy change for the hydrogenation of ethene



Energy required to break bonds: $4 \times (\text{C–H}) + (\text{C=C}) + (\text{H–H}) = 2700 \text{ kJ mol}^{-1}$
 Energy released in forming new bonds: $6 \times (\text{C–H}) + 1 \times (\text{C–C}) = 2825 \text{ kJ mol}^{-1}$
 Enthalpy change: $2700 - 2825 = -125 \text{ kJ mol}^{-1}$

b) the enthalpy change for the cracking of decane



Energy required to break bonds: $22 \times (\text{C–H}) + 9 \times (\text{C–C}) = 12\,209 \text{ kJ mol}^{-1}$
 Energy released in forming new bonds: $22 \times (\text{C–H}) + (\text{C=C}) + 7 \times (\text{C–C}) = 12\,127 \text{ kJ mol}^{-1}$
 Enthalpy change: $12\,127 - 12\,209 = +82 \text{ kJ mol}^{-1}$
 N.B. This can be more easily calculated by realising that the overall reaction involves breaking of 2 C–C bonds and formation of 1 new C=C bond. The calculation is then
 Energy required to break bonds: $2 \times (\text{C–C}) = 694 \text{ kJ mol}^{-1}$
 Energy released in forming new bonds: $1 \times (\text{C=C}) = 612 \text{ kJ mol}^{-1}$
 Enthalpy change: $694 - 612 = +82 \text{ kJ mol}^{-1}$

12. Explain in terms of bond breaking and bond formation why combustion reactions are exothermic but cracking reactions are endothermic.

In combustion reactions, the bond enthalpies in the products are greater than those in the reactants. Therefore, more energy is released in making the bonds in the products than is required to break the bonds in the reactants.

In cracking reactions, the bond enthalpies in the products are smaller than those in the reactants. Therefore, less energy is released in making the bonds in the products than is required to break the bonds in the reactants.

CHEMISTRY A AND CHEMISTRY B (SALTERS)

Learner Activity

OCR Resources: *the small print*

OCR's resources are provided to support the teaching of OCR specifications, but in no way constitute an endorsed teaching method that is required by the Board, and the decision to use them lies with the individual teacher. Whilst every effort is made to ensure the accuracy of the content, OCR cannot be held responsible for any errors or omissions within these resources.

© OCR 2017 - This resource may be freely copied and distributed, as long as the OCR logo and this message remain intact and OCR is acknowledged as the originator of this work.

OCR acknowledges the use of the following content: n/a

Please get in touch if you want to discuss the accessibility of resources we offer to support delivery of our qualifications: resources.feedback@ocr.org.uk